

The Mathematics of Radioactive Decay

1) Discovery of the radioactive decay law. In 1900, Ernest Rutherford noticed a decrease over time in the radioactive intensity of a sample he was studying, so he began to measure this decrease. He determined that the decrease fit the following formula (using modern symbols):

$$N = N_0 e^{-kt} \quad (1)$$

where N = the amount of sample remaining after an amount of time 't' and N_0 = the original starting amount of the sample. e has its usual meaning and k is a constant with a unique value for each substance. Rutherford named it the radioactive decay constant.

2) Straight-line form of the decay law. Equation 1 can be modified to the form of a straight line formula ($y = mx + b$) as follows:

$$\begin{aligned} N / N_0 &= e^{-kt} && \text{by rearrangement} \\ \ln(N / N_0) &= -kt && \text{take natural logarithm of each side} \quad (2) \\ \ln N - \ln N_0 &= -kt && \text{division of exponents is done by subtraction} \\ \ln N &= -kt + \ln N_0 && \text{by rearrangement} \end{aligned}$$

This last equation fits the general form of a straight-line equation. The y-axis is the natural log of N and the x-axis is t (time). The line will run from upper left to lower right (negative slope) with negative k being the slope.

3) An equation for calculating half-life. The equation (2) can be modified to give an equation where knowing the value for k leads directly to the length of the half-life.

$$t = \frac{-\ln(N / N_0)}{k} \quad \text{by rearrangement}$$

From the definition of half-life, we obtain the value for N / N_0 as one-half. After one half-life, half the material on-hand at the start will have decayed. If the starting amount equals 2, then the ending amount one half-life later will equal one.

Considering the numerator only: $-\ln(1/2) = -(\ln 1 - \ln 2) = -(0 - \ln 2) = \ln 2$

Letting $T_{1/2}$ = time of one half-life, we obtain:

$$T_{1/2} = 0.693 / k \quad \text{since } \ln 2 = 0.693$$

4) Miscellaneous half-life information. The fraction one-half figures prominently in this.

$$(1/2)^n \quad \text{where } n = \text{the number of half-lives yields the decimal portion of substance remaining.}$$

If the decimal fraction of substance remaining is known, setting it equal to $(1/2)^n$ and solving will yield the number of half-lives elapsed.